

IMPACT OF **CLEAN DIESEL** TECHNOLOGY ON CLIMATE CHANGE


*Robert McGraw and James Wegrzyn
Brookhaven National Laboratory
Upton, NY 11973*

Overview:

- **Survey of emissions and BC aerosol processes**
- **New developments in modeling (aerosols, clouds, and climate)**
- **Future studies**

AEROSOL PROCESSES AND THEIR IMPACT ON CLIMATE

Understanding and modeling the importance of diesel emissions reduction on climate



SOME KEY QUESTIONS:

Emissions: What is the contribution of diesel as a primary BC aerosol source?

(including impact of clean diesel technology and lower emissions of CO₂ on climate)


What is the subsequent mixing state of the BC aerosol? (e.g. does BC become coated with sulfate?)

Is mixing state important to the assessment of BC aerosol optical properties, cloud properties, and climate effects?


THE UNCERTAIN CONTRIBUTION OF AEROSOLS TO CLIMATE

RADIATIVE FORCING OVER THE INDUSTRIAL PERIOD IPCC (2001)

GHG's and aerosol direct and indirect effects



Summary for Policymakers A Report of Working Group I of the Intergovernmental Panel on Climate Change



EMISSIONS





Figure 2. Emissions of black carbon aerosol. (a) “Contained” combustion, based on 1996 activity data. (b) Open burning, annual average. The color coding is an approximately logarithmic scale. Units are $\text{ng}/\text{m}^2/\text{s}$ ($1 \text{ ng}/\text{m}^2/\text{s} \sim 32 \text{ kg}/\text{km}^2/\text{yr}$).

T. Bond et al., “A technology-based inventory of black carbon and organic carbon emissions from combustion.”
J. Geophys. Res. 109, D14203 (2004)

EMISSIONS (CONTINUED)



From Bond et al. J. Geophys. Res. 109, D14203 (2004)

DIRECT EFFECTS OF BLACK CARBON AEROSOLS ON CLIMATE



Figure from Andreae M. O. “The dark side of aerosols”, *Nature* 409, 671-672 (2001).

Importance of mixing state: Sulfate coated-BC can have double the contribution to global warming relative to the uncoated form (Jacobson, 2002) → Greater contribution to warming than previously thought
→ Greater beneficial impact from BC reduction than previously thought.

Optical absorption properties: these were based on calculations for a spherical BC core/sulfate mantle particle. Observed particles have more complicated shapes.

AEROSOL INDIRECT EFFECTS ON CLIMATE

Indirect effects: **Aerosols** → **Clouds** → **Climate**


- Aerosols, by providing more seed particles for cloud condensation, partition cloud water into more droplets of smaller size, resulting in optically thicker and more reflecting clouds (Twomey effect or first indirect effect).
- This same partitioning into smaller drops leads to a decrease in precipitation resulting in longer cloud lifetimes and greater cloud cover (second indirect effect). *Clouds tend to persist longer under polluted conditions than in the marine environment.*
- However, the issue is further complicated with BC due to absorption: By warming cloudy air, BC can result in enhanced evaporation resulting in dissipation or partial dissipation of clouds (semi-direct effect).

ABSORBING PHENOMENA



Aircraft photo taken during INDOEX 1999 over the tropical Indian Ocean shows small cumulus clouds embedded in an absorbing aerosol layer. Absorption of solar radiation by the aerosol heats the air surrounding the clouds, causing them to evaporate. (*From Schwartz and Buseck, Science 288, 989 (2000)*).

IMPACT ASSESSMENT CONSIDERATIONS



10-nm soot carbon spherules can be found embedded within much larger sulfate particles.
Figure from Buseck and Posfai (1999).

The inhomogeneity in properties and geographical distributions of BC aerosols make it difficult to characterize their influences on climate and to represent these influences in models.

Our approach: Use a statistically-based representation of generally-mixed aerosols, and their physical and optical properties, suitable for use in regional to hemispheric scale atmospheric models.

**NEW DEVELOPMENTS IN THE MODELING OF
GENERALLY-MIXED PARTICLE POPULATIONS
USING THE METHOD OF MOMENTS**

MOMENTS OF THE PARTICLE SIZE DISTRIBUTION

$$\mu_k \equiv \int_0^{\infty} r^k \left(\frac{dN}{dr}\right) dr \quad (\text{not normalized})$$

Moment	Physical Interpretation	Unit
μ_0	Particle number concentration	cm^{-3}
μ_1	Total radius per unit volume	cm cm^{-3}
μ_2	$(4\pi)^{-1} \times$ Area per unit volume	$\text{cm}^2 \text{cm}^{-3}$
μ_3	$\left(\frac{4\pi}{3}\right)^{-1} \times$ Volume per unit volume	$\text{cm}^3 \text{cm}^{-3}$

QUADRATURE METHOD OF MOMENTS

The *quadrature* method of moments evaluates the moment evolution equation

$$\frac{d}{dt}\mu_k = k \int_0^{\infty} r^{k-1} \phi(r) f(r) dr$$

by Gaussian quadratures:

$$\frac{d}{dt}\mu_k \cong k \sum_{i=1}^3 r_i^{k-1} \phi(r_i) w_i$$



This approach is *completely general and highly accurate*.

R. McGraw, *Aerosol Sci. Technol.* (1997)

INCORPORATION OF QMOM IN REGIONAL SCALE MODEL

Host model: MAQSIP - Multiscale Air Quality Simulation Platform - prototype of US EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System

Model domain: 72 by 74 square, 36-km grid cells, 22 layers, sigma coordinate, from the surface to ~160 hPa.

Meteorological driver: MM5 meteorological model.

Emissions: 1990 EPA National Emissions Trends (NET90) Inventory.

Chemistry: SO₂ oxidation by OH in air and oxidation by H₂O₂ in cloud droplets.


Clouds: Sub-grid convective (precipitating and non-precipitating) and grid-scale resolved.


Particle activation, wet and dry deposition: Dependent on particle size.

Moment-Based Simulation of Microphysical Properties of Sulfate Aerosols in the Eastern United States

Shaocai Yu, Prasad S. Kasibhatla, Douglas L. Wright, Stephen E. Schwartz, Robert McGraw, Aijun Deng, J. Geophysical Res. 108, D12, 4353 (2003).

Distributions of mean sulfate mixing ratios and number concentrations.









Predicted aerosol size distributions from moments and comparison with observations

OPTICAL PROPERTIES

Results, illustrated here for forcing and 28 test distribution, obtainable directly from six moments



Squares: exact results calculated from size distribution


Circles: Results obtained from six radial moments

BIVARIATE EXTENSION OF THE QMOM FOR MODELING SIMULTANEOUS COAGULATION AND SINTERING OF PARTICLE POPULATIONS


Wright, McGraw, and Rosner, *J. Coll. Interface Sci.* 236, 242 (2001)

COALESCENCE
RESTRUCTURING
↑
FRACTAL AGGREGATES
↑
COAGULATION
↑
SPHERULE PRODUCTION
↑
COMBUSTION

initial distribution (log scale)



final distribution (2D sectional model)




Compared direct moment evolution using the QMOM with results from a 2D (150x150) discrete model for 36 mixed moments

- max error = 7%
- cpu time:

12 days

(sectional)

10 s (QMOM)



NEW DEVELOPMENTS IN THE MODELING OF GENERALLY-MIXED AEROSOLS



Aerosol Science 35 (2004) 561–576

*Journal of
Aerosol Science*

www.elsevier.com/locate/jaerosci

Representation of generally mixed multivariate aerosols by the
quadrature method of moments: I. Statistical foundation

Choongseok Yoon^{a,b,1}, Robert McGraw^{a,*}

^a*Atmospheric Sciences Division, Brookhaven National Laboratory, Upton, NY 11973, USA*

^b*Department of Applied Mathematics and Statistics, State University of New York,
Stony Brook, NY 11794-3600, USA*


Aerosol Science 35 (2004) 577–598

Representation of generally mixed multivariate aerosols by the
quadrature method of moments: II. Aerosol dynamics

AEROSOLS, CLOUDS, AND PRECIPITATION

New model for drizzle formation, R. McGraw and Y. Liu

- *Phys. Rev. Letts.* **90**, 018501, 1-4 (2003)
- *Geophys. Res. Letts.* **31**, L06121, doi: 10.1029/2003GL019117 (2004)
- *Phys. Rev. E.*, in press (2004)



*Contours of constant drizzle rate (solid)
and average cloud droplet radius (dashed)*

N_D = droplet number concentration
L = liquid water fraction

DRIZZLE THRESHOLD

$$\log_{10} J_{SS} (\text{cm}^3 \text{s}^{-1}) = -6 \text{ to } -5$$

(1-10 drops/m³ s)

(region within the two lowest
solid contours)

FUTURE STUDIES

- **Simulation of BC aerosol and cloud processes using a multi-scale (regional to hemispheric scale) chemical transport model.**
- **These models will include the latest inventories for diesel BC emissions and projected emissions reduction achievable through clean diesel technology. The objective will be to quantify the favorable climate impact now achievable through BC particulate emissions reduction**