

Robust Non-Oxidative Glycolysis in *Escherichia coli*

Paul P. Lin¹ (phlin@ucla.edu), Alec J. Jaeger,¹ Maria Chun,¹ Kouki M. Yoshino,¹ Sawako Konishi,¹ Sharon C. Xu,¹ and **James C. Liao**¹

¹Department of Chemical and Biomolecular Engineering, University of California, Los Angeles

Project Goals: This project seeks to construct strains of *Escherichia coli* to rely solely on Non-oxidative glycolysis (NOG) for glucose catabolism during aerobic growth, fundamentally rewiring the central metabolism to favor carbon conservation in acetyl-CoA biosynthesis. This work will also engineer the constructed strains for energy applications. Specifically, we aim to produce an advanced biofuel, n-butanol, with close to complete carbon conservation. Hydrogen or formate will be provided to supply reducing equivalents needed to drive the NADH dependent pathway.

Acetyl-coenzyme A (acetyl-CoA) is a two carbon metabolite and important metabolic precursor to a variety of industrially relevant compounds including biofuels. An ultimate limitation of acetyl-CoA derived biochemical production is the inherent carbon loss when forming acetyl-CoA. Most organisms use some glycolytic variation, commonly the Embden-Meyerhof Pathway (EMP), to initially degrade sugar into pyruvate. Pyruvate, a C₃ metabolite, is then decarboxylated to form acetyl-CoA, losing carbon to the environment. This decarboxylation limits the carbon yield to only two molecules of acetyl-CoA from one molecule of hexose, thus inhibiting the economics of any associated bioprocess. A synthetic sugar catabolism pathway, termed non-oxidative glycolysis (NOG), was recently developed to address this problem, as it uses a combination of phosphoketolase dependent cleavage of sugar phosphates and a carbon rearrangement cycle to directly generate three C₂ units per hexose in a redox neutral manner. To further expand the applications using NOG, an *Escherichia coli* strain was constructed to rely solely on NOG for sugar catabolism in this work. Therefore, the resulting strain offers significant potential to be engineered for the production of a variety of acetyl-CoA derived compounds. To implement NOG as a growth pathway, all native sugar degradation pathways, including the EMP, ED and methylglyoxal bypass, were removed, eliminating the cell's ability to grow on sugar as a sole carbon source. In addition, the glyoxylate shunt and gluconeogenesis pathways, which are necessary for the production of essential metabolites using NOG, were upregulated. Then, this engineered strain was evolved to grow in minimal glucose media supplemented with exogenous acetate. The overexpression of critical NOG protein phosphoketolase restored the cells ability to grow on sugar as a sole carbon source and further evolution improved the cell's growth phenotype in glucose and xylose minimal media. Under anaerobic conditions, it was verified that this strain produces acetate as a major fermentation product from glucose. The carbon rearrangement pathway will be further overexpressed to increase the carbon yield through NOG pathway for acetate production. To further demonstrate the potential of NOG, this strain will be additionally engineered to produce C₂ derived biofuels by supplying reducing power in the form of hydrogen or formate.